

MICROVUE™

Bone Health

Technical Data Sheet

MicroVueTRAP5b

For **Research Use Only**. Not for use in diagnostic procedures.

The MicroVue TRAP5b assay is an enzyme immunoassay for the quantitative determination of tartrate-resistant acid phosphatase isoform 5b (TRAcP 5b) in Human Serum and Plasma.

Format

- ELISA
- 96-well assay plate
- Sample type: EDTA or Heparin Plasma and Serum
- Controls: High, Low

Species Reactivity

- Reactive with human derived TRAP5b in serum, plasma or cell-culture medium.
- No reactivity detected in Baboon, Cow, Chicken, Cynomologous Macaque, Goat, Mouse, Rabbit, Rat, Rhesus Macaque, and Sheep.

Assay Steps

- Add 100 µL of Sample Diluent.
- Add 50 µL of Standards, Controls and specimens.
- Seal the microplate. Incubate for 60 minutes at room temperature on a microplate shaker.
- Wash the microplate wells three times with Wash Buffer.
- Add 100 µL of Substrate Solution.
- Seal the microplate. Incubate for 60 minutes at 37°C.
- Add 50 µL Stop Solution and read the OD at 405 nm.
- Data reduction: Quadratic.

Performance Characteristics

Method: ELISA

Analyte: Tartrate-resistant acid phosphatase isoform 5b (TRAcP5b)

Specimen Volume: 50 µL

Sensitivity: 0.2 U/L

Assay Range: 0.2-15 U/L

Precision CVs within-run: 3-8%

Precision CVs between-run: 4-8%

Incubation Time: 2 Hours

TO ORDER: Call Quidel Customer service at 1.800.524.6318 (USA only), 1.408.616.4301, or fax 1.408.616.4310.

MicroVue TRAP5b

Cat. #8033



RESEARCH TO RAPIDS™

Worldwide Headquarters

10165 McKellar Court
San Diego, CA 92121 USA
www.quidel.com

For more information on our products and other Quidel locations, visit our website.
800.524.6318 • 408.616.4301 • 408.616.4311 (fax)

0133BH0206D-3 (01/10)