



Trypsin Solution

Additive when preparing cell culture media.

FOR *IN VITRO* DIAGNOSTIC USE

DESCRIPTION

One bottle containing 7 mL of a 25 mg/mL trypsin solution (porcine origin) in 0.9% saline for use as an additive when preparing cell culturing media. Trypsin is packaged and shipped frozen.

WARNINGS AND PRECAUTIONS

- General purpose laboratory reagent.
- Use undiluted.
- May cause allergic skin reaction.
- Testing should be performed in an area with adequate ventilation.
- Dispose of containers and unused contents in accordance with Federal, State and Local regulatory requirements.
- Wear suitable protective clothing, gloves, and eye/face protection when handling the contents of this kit.
- Wash hands thoroughly after handling.
- For additional information on hazard symbols, safety, handling and disposal of the components within this kit, please refer to the Safety Data Sheet (SDS) located at quidel.com.

STORAGE

Reagent will retain its full potency through the expiration date when stored at -20°C .

ASSISTANCE

To place an order or for technical support, please contact a Quidel Representative at 800.874.1517 (in the U.S.) or 858.552.1100 (outside the U.S.), Monday through Friday, from 8:00 a.m. to 5:00 p.m., Eastern Time. Orders may also be placed by fax at (740) 592-9820. For e-mail support contact customerservice@quidel.com or technicalsupport@quidel.com.

For services outside the U.S., please contact your local distributor. Additional information about Quidel, our products, and our distributors can be found on our website quidel.com.

REF

10-110007 – Trypsin Solution

IVD



EC REP

MDSS GmbH
Schiffgraben 41
30175 Hannover,
Germany



Diagnostic Hybrids, Inc. – a subsidiary of Quidel Corporation
2005 East State Street, Suite 100
Athens, OH 45701 USA
quidel.com

PI9041000EN00 (09/18)

REF

Catalogue number



CE mark of conformity

EC REP

Authorized Representative
in the European Community

LOT

Batch code



Use by



Manufacturer



Temperature limitation



Intended use



Consult e-labeling
instructions for use



Do not reuse

IVD

For *In Vitro* diagnostic use

CONT

Contents/Contains
